

STIC-ILL

From: Nguyen, Bao-Thuy  
Sent: Wednesday, September 17, 2003 1:03 PM  
To: STIC-ILL  
Subject: Reference - 09/857,213

*PH442.643*  
*mu*  
*Adams*

Please give me a copy of the following references ASAP:

1. Kwak et al., Gene, 1996. Vol. 173(2), 257-259.
2. Giunta et al., Annals of the New York Academy of Sciences. 1992, 673, 342-349.
3. Ayrault-Jarrier et al., Advances in Experimental Medicine and Biology. 1988, 243, 149-155.
4. Fidge et al., Biochimica et Biophysica Acta. 1989, 1003(1), 84-90.
5. Milthorp et al., Arteriosclerosis, 1986, 6(3), 295-296.

Bao-Thuy Nguyen  
(703) 308-4243  
CM 1, Room 7E05

GENE 09675

## Cloning and characterization of cDNAs coding for heavy and light chains of a monoclonal antibody (MabA34) specific for human plasma apolipoprotein A-I

(Recombinant DNA; immunoglobulin Fab; Apo A-I; polymerase chain reaction; high-density lipoproteins; HDL cholesterol; DNA sequencing)

Ju-Won Kwak<sup>a</sup>, Dong-Ik Lee<sup>a</sup>, Byung-Kwon Choi<sup>b</sup>, Won-Kyung Cho<sup>a</sup>, Sang-Han Lee<sup>c</sup>,  
Yong-Bok Park<sup>d</sup> and Moon Hi Han<sup>a</sup>

<sup>a</sup>Korea Research Institute of Bioscience and Biotechnology, KIST, Daeduck Science Town, Taejeon 305-600, South Korea; <sup>b</sup>Sam-Il Pharmaceutical Co. Ltd., Banghae-Dong, Seocho-Ku, Seoul 137-060, South Korea; <sup>c</sup>Biochemistry Laboratory, College of Medicine, Soonchunhyang University, Chonan 330-020, South Korea; and <sup>d</sup>Department of Genetic Engineering, Kyungpook National University, Taegu 702-701, South Korea

Received by H.G. Zachau: 2 November 1995; Revised/Accepted: 4 December 1995; Received at publishers: 26 January 1996

BEST AVAILABLE COPY

### SUMMARY

We have determined the nucleotide (nt) sequences encoding the heavy (H)- and light (L)-chains of the Fab fragment of a murine monoclonal antibody, MabA34 ( $\gamma$ 1,  $\kappa$ ), which is specific for human plasma apolipoprotein A-I of high-density lipoproteins. The variable (V) regions of the H- and L-chains were revealed to be members of mouse H-chain subgroup II(A) and  $\kappa$  L-chain subgroup II, respectively. A few unusual amino acids in the V region of the H-chain and nt residues probably introduced by somatic mutations from germline genes were also identified.

### INTRODUCTION

Plasma apolipoprotein (Apo) A-I of high-density lipoproteins (HDL) has received considerable clinical attention because the plasma concentrations have an inverse relationship with the incidence of coronary artery disease (Avogaro et al., 1979; Naito, 1986; Bachorik and

Kwiterovich, 1988). Apo A-I is a major protein component of HDL which is mainly responsible for the reverse cholesterol transport from peripheral tissues to the liver. Following the recognition of an inverse correlation between plasma HDL and the incidence of coronary artery disease, Apo A-I has become the subject for numerous immunochemical studies involving both polyclonal and monoclonal Ab. Here, we report the cloning and characterization of cDNAs encoding the heavy (H)- and light (L)-chains for the Fab fragment of a mAb (MabA34) which specifically recognizes and binds to human plasma Apo A-I of HDL.

Correspondence to: Dr. J.-W. Kwak, Protein Engineering Research Group, Korea Research Institute of Bioscience and Biotechnology, KIST, P.O. Box 115, Yuseong, Taejeon 305-600, South Korea. Tel. (82-42) 860-4125; Fax (82-42) 860-4594; e-mail: jwkwak@geri4680.geri.re.kr

Abbreviations: aa, amino acid(s); Ab, antibody(ies); Apo, apolipoprotein(s); bp, base pair(s); C, constant; cDNA, complementary DNA; CDR, complementarity determining region(s); C<sub>H</sub>1, H-chain C region 1; C<sub>L</sub>,  $\kappa$  L-chain C region; D, diversity; Fab, antigen-binding Ab fragment; FR, framework region(s); H, heavy; HDL, high-density lipoprotein(s); J<sub>H</sub>, H-chain joining segment; J<sub>L</sub>,  $\kappa$  L-chain joining segment; L, light; mAb, monoclonal Ab; nt, nucleotide(s); oligo, oligodeoxyribonucleotide; PCR, polymerase chain reaction; V, variable; V<sub>H</sub>, H-chain V region; V<sub>L</sub>, L-chain V region.

### EXPERIMENTAL AND DISCUSSION

#### (a) Cloning of MabA34-specific heavy and light chain cDNAs

The fusion of mouse Balb/c spleen cells and Sp2/O-Ag-14 myeloma cells has produced the hybridoma cell

Asp<sup>26</sup>. Otherwise, all the invariant aa for the murine H-chain subgroup II(A) were conserved (Harris and Bajorath, 1995). A total of five Cys in the H-chain Fab region and five Cys in the L-chain were found, which would be involved in intra- or inter-chain disulfide bonding necessary to form the Fab immunoglobulin structure. At the end of the H-chain V region (aa 100–112 in Fig. 1), J<sub>H</sub>2 element without any nt mismatch was identified. In the CDR3 of the H-chain, a unique D region comprised of only one aa (Pro<sup>99</sup>) was also identified. Comparisons with germline V genes have assigned the V region of H-chain to 19.1.2, a VhJ558 family (Akolkar et al., 1987), and suggested that the V region had undergone somatic mutations characteristic of an antigen-driven immune response.

The sequence analysis of L-chain-specific cDNA of MabA34 revealed that the V region of the L-chain is a member of mouse  $\kappa$  chain subgroup II (Fig. 2). All the invariant aa for the  $\kappa$  chain were conserved with no exceptions. The CDR 1, 2, and 3 were positioned at aa 24–39/55–61/94–102, respectively. At the end of L-chain V region (aa 100–111 in Fig. 2), J<sub>H</sub> element with 1 nt mismatch (GGG instead of TGG for Arg<sup>100</sup>) was identified. Comparisons with germline V genes allowed to determine K1A5 (Corbet et al., 1987) as a likely germline progenitor of the V domain of the L-chain.

[illegible]

**Fig. 2.** Nucleotide sequence of the cDNA coding for the L-chain of monoclonal antibody MabA34 specific for human plasma Apo A-I, and deduced aa sequence. The  $V_L$  and  $C_L$  correspond to aa 1-112 and aa 113-219, respectively. The regions of CDR 1, 2, 3 are blocked. The progenitory, mouse germline V gene sequence of KIA5 (Corbet et al., 1987) was aligned with the sequence to show the nt residues probably introduced by somatic mutations. The dashes denote identity. The sequence is available from GenBank database under accession No. U29147. **Methods:** The PCR product subcloned into pBluescript (Stratagene, La Jolla, CA, USA) was fully sequenced on both orientations.

The sequence analyses suggest that the H- and L-chain-specific cDNAs are functional. Comparison of nt sequences of the H- and L-chain V region with the available immunoglobulin genes listed in GenBank database showed that the sequences have not been previously reported. The cDNAs will be used for expression and serum-free production in *Escherichia coli* of the recombinant Ab, which is valuable for clinical diagnostics as well as for numerous immunochemical studies of human plasma Apo A-I.

### (c) Conclusions

We have cloned and determined the nt sequences of the heavy- and light-chain-specific mAb-encoding cDNAs of murine MabA34 ( $\gamma 1$ ,  $\kappa$ ) specific for human plasma Apo A-I of HDL. The sequence analyses revealed that both cDNAs are functional, and the variable regions of the heavy and light chains are members of mouse H-chain subgroup II(A) and  $\kappa$  L-chain subgroup II, respectively. These sequences have not been previously reported.

### ACKNOWLEDGEMENTS

This work was supported by a G7 research grant to J.W.K. from the Ministry of Science and Technology, Korea.

### REFERENCES

- Akolkar, P.N., Sikder, S.K., Bhattacharya, S.B., Liao, J., Gruezo, F., Morrison, S.L. and Kabat, E.A.: Different  $V_L$  and  $V_H$  germ-line genes are used to produce similar combining sites with specificity for  $\alpha(1 \rightarrow 6)$  dextrans. *J. Immunol.* 138 (1987) 4472-4479.
- Avogaro, P., Bittolo Bon G., Cazzolato, G. and Quinci, G.B.: Are apolipoproteins better discriminators than lipids for atherosclerosis? *Lancet* 28 (1979) 901-903.
- Bachorik, P.S. and Kwiterovich Jr., P.O.: Apolipoprotein measurements in clinical biochemistry and their utility vis-a-vis conventional assays. *Clin. Chim. Acta* 178 (1988) 1-34.
- Corbet, S., Milili, M., Fougereau, M. and Schiff, C.: Two  $V_L$  germ-line genes related to the GAT idiotype network (Ab1 and Ab3/Ab1') account for the major subfamilies of the mouse  $V_L$ -1 variability subgroup. *J. Immunol.* 138 (1987) 932-939.
- Harris, L. and Bajorath, J.: Profiles for the analysis of immunoglobulin sequences: comparison of V gene subgroups. *Protein Sci.* 4 (1995) 306-310.
- Kabat, E.A., Wu, T.T., Reid-Miller, M., Perry, H.M., Gottesman, K.S. and Foeller, C.: Sequence of Proteins of Immunological Interest, 5th ed. US Department of Health and Human Services, Public Health Services, National Institute of Health, Bethesda, MD, 1991.
- Naito, J.K.: The clinical significance of apolipoprotein measurements. *J. Clin. Immunoassay* 9 (1986) 11-20.
- Sanger, F., Nicklen, S. and Coulson, A.R.: DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74 (1977) 5463-5467.

BEST AVAILABLE COPY